

- 19 Pugh LG, Corbett JL, Johnson RH. Rectal temperatures, weight losses, and sweat rates in marathon running. *J Appl Physiol* 1967;**23**:347–52.
- 20 Noakes TD, Carter JW. Biochemical parameters in athletes before and after having run 160 kilometres. *S Afr Med J* 1976;**50**:1562–6.
- 21 Cohen I, Zimmerman AL. Changes in serum electrolyte levels during marathon running. *S Afr Med J* 1978;**53**:449–53.

Exercise-associated hyponatraemia: facts and myths

In an elaborate manuscript, Noakes and Speedy argue that exercise-associated hyponatraemia (EAH) is due to overhydration and that the usual sodium deficit incurred during exercise plays an insignificant role.^{1,2} Furthermore, these researchers claim that the reluctance to accept their evidence may be because it conflicts with the prevalent message of the sports drink industry. This approach is, however, oversimplification of the situation and at best singlet-sided. Noakes and Speedy in actuality prove the obvious. Overhydration will dilute body fluid, and thus plasma sodium concentration, as the primary extracellular electrolyte, will consequently be reduced. However, because hyponatraemia can be observed in the presence of a normal, depleted or expanded plasma volume, the issue of hyponatraemia should be approached from a broader perspective. We emphasise that the inability to replace cumulative sodium losses in sweat during long physical exertions is not less important as the background for EAH, whereas hyperhydration is only a marginal, although vastly reported, condition associated with EAH.

Body fluid homeostasis is a primary variable, which is important in the stability of the “inner environment”. Therefore, total body water and the proportion of extracellular to intracellular volumes should be maintained very delicately. Dehydration during exercise results in graded physiological responses, including a decrease in cardiac output despite an increase in heart rate, an increase in core temperature and a decrease in peripheral blood flow, which ultimately affects performance. Eichna *et al*³ described the ability of dehydrated participants to perform in the heat as: “total incapacitation ... acclimatised subjects who had performed a given task easily ... reduced to apathetic, listless, plodding men straining to finish the same task”. Fluid replacement during exercise contributes to the maintenance of plasma volume, aiding thermal and cardiovascular homeostasis and thus reversing all these adverse effects.^{4,5} Therefore, dehydration should not be underestimated: it will not only adversely influence performance but will also, when severe, threaten life. Sport specialists who advocate fluid replacement *ad libitum* actually disregard the ill effects of dehydration. However, optimal rehydration should not be withheld only out of fear for overhydration.

To illustrate the point that overhydration is not the sole reason for EAH, we analyse the following examples: Hew-Butler *et al*⁶ show that following the South African Ironman triathlon (lasting 12:20 h), serum $[Na^+]$ was maintained at 140.9 mEq/L (baseline levels were 140.5 mEq/L). This was achieved by drinking *ad libitum*, thus attaining a dehydration level of 3.9%. If the athletes would have drunk water according to sweat loss, with no overhydration at all, final serum $[Na^+]$ would have been in the magnitude of 134.7 mEq/L (assuming a sweat rate of 450 ml/h and a sweat sodium concentration of 50 mEq/L). Similarly, Montain *et al*⁷ show that serum $[Na^+]$ of a 70 kg male athlete after a 9 h, 90 km

ultramarathon foot race will be 130.2 mEq/L if $[Na^+]$ in sweat is 50 mEq/L and if he drinks water to maintain euhydration. By recalculating the data, assuming that the athlete would have drunk *ad libitum*, incurring a mild state of dehydration at an estimate level of 4%, serum $[Na^+]$ after the race will be higher, 135.5 mEq/L. These two examples well illustrate that during long physical exertions, overhydration is not the reason for lowering serum $[Na^+]$ but rather the cumulative loss of sodium in sweat that is not adequately replenished. This opinion is substantiated in the following observation by Y Epstein. Two groups of soldiers participated in a 50 km (10 hr) intense foot march. Group A drank water at a rate that compensated sweat loss (1000 ml/h). Group B drank the same amount of water but were also served with snacks and sandwiches, which they consumed *ad libitum* (average total sodium consumption = 222 mEq). At the end of the exercise, the mean (SD) serum $[Na^+]$ of participants in group A was 134 (1) mEq/L, whereas in group B it was 138 (1) mEq/L (pre-exercise $[Na^+] = 142$ mEq/L). The consumption of sodium by participants in group B has attenuated the drop in plasma $[Na^+]$. This finding is corroborated by Vrijens and Rehrer, who showed that sodium consumption can slow down the drop in plasma $[Na^+]$ after long physical exercise.⁸ The small reduction in serum $[Na^+]$ in group B is due to the less than desirable sodium intake. Assuming a sweat $[Na^+]$ of 40–50 mEq/L, these soldiers replaced only about 50% of the amount of sodium loss.

Is salt replacement needed all the time? The answer is probably “no”. Sweat is a hypotonic fluid and the reported $[Na^+]$ in sweat is in the range of 20–80 mEq/L.⁹ In general, $[Na^+]$ in the sweat of fit acclimated subjects is in the range of 20–40 mEq/L, whereas in unfit subjects it is nearly twofold higher.¹⁰ Thus, when water intake is inadequate to compensate sweat loss and dehydration ensues, the blood becomes hypertonic. When water intake is equal to water loss by sweating, theoretically the plasma electrolyte levels, especially that of sodium, should fall. However, owing to the high plasma sodium content and the large extracellular fluid volume, hyponatraemia ($[Na^+] < 135$ mEq/L) is not apparent until late in exercise, after large amounts of water have been ingested with no sodium replacement.

Hyponatraemia is seldom expected in athletes exercising at high intensities for periods of <4 h.^{7,11} This conclusion is based on the estimated low total loss of sodium in sweat and the general observation that at high work rates, fluid intake is less than optimal to maintain euhydration. Under these conditions, the prime concern is replenishing body fluid by an adequate drinking routine. During longer periods of exercise, hypovolaemia and dehydration are expected, but hyponatraemia, due to cumulative sodium loss, may also develop;¹¹ the above examples prove this. It should, however, be noted that slow, unfit athletes are at a higher risk of developing EAH owing to overt overhydration, in the face of their lower sweat rate and relatively large sodium losses in sweat, as recently shown by Almond *et al*¹² during the Boston marathon.

In conclusion, to maintain euvolaemia and serum osmolality, water and sodium intake must compensate for sweat losses. If fluid consumption exceeds sweat rate, dilutional hyponatraemia may occur. As sodium concentration in sweat is low, especially in trained athletes, salt consumption is not required for exercises of short duration (<4 h). However, during prolonged exercise, hyponatraemia will

develop even without excess fluid intake, unless sweat sodium loss is adequately replaced. The most effective and physiological method of replacing sodium losses is not necessarily by consuming sports drinks, but rather by appropriate diet. This is the common practice in Israel and the attitude recommended by the American College of Sports Medicine (ACSM).¹³ Such a routine prevents dehydration and discourages overhydration, and also prevents the development of hyponatraemia.

Yoram Epstein, Yoav Cohen-Sivan

Sheba Medical Center, Heller Institute of Medical Research, Tel-Hashomer, Israel

Correspondence to: Professor Y Epstein, Sheba Medical Center, Heller Institute of Medical Research, Tel-Hashomer 52621, Israel; yoram.epstein@sheba.health.gov.il

Competing interests: None declared.

References

- 1 Noakes TD, Speedy DB. Case proven: exercise associated hyponatremia is due to overdrinking. So why did it take 20 years before the original evidence was accepted? *Br J Sports Med* 2000;**40**:567–72.
- 2 Noakes TD, Sharwood K, Perkins DR. The dipsomania of great distance: water intoxication in an Ironman triathlete. *Br J Sports Med* 2004;**38**:e16.
- 3 Eichna LW, Bean WB, Ashe WF, *et al*. Performance in relation to environmental temperature. Reaction of normal young men to hot, humid (simulated jungle) conditions. *Bull Johns Hopkins Hosp* 1945;**76**:25–58.
- 4 Sawka MN, Pandolf KB. Effects of body water loss on physiological function and exercise performance. Gisolfi CV, Lamb DR, eds. *Perspective in exercise science and sports medicine. Vol 3: fluid homeostasis during exercise*. Carmel, IN: Benchmark Press, 1990:1–30.
- 5 Montain SJ, Coyle EF. Influence of graded dehydration on hyperthermia and cardiovascular drift during exercise. *J Appl Physiol* 1992;**73**:1340–50.
- 6 Hew-Butler TD, Sharwood K, Collins M, *et al*. Sodium supplementation is not required to maintain serum sodium concentrations during Ironman triathlon. *Br J Sports Med* 2006;**40**:255–9.
- 7 Montain SJ, Sawka MN, Wenger CB. Hyponatremia associated with exercise: risk factors and pathogenesis. *Exerc Sports Sci Rev* 2001;**29**:113–7.
- 8 Vrijens DM, Rehrer NJ. Sodium-free fluid ingestion decreases plasma sodium during exercise in the heat. *J Appl Physiol* 1999;**86**:1847–51.
- 9 Rehrer NJ. Fluid and electrolyte balance in ultra-endurance sport. *Sports Med* 2001;**31**:701–15.
- 10 Allan JR, Wilson CG. Influence of acclimatization on sweat sodium concentration. *J Appl Physiol* 1971;**30**:708–12.
- 11 Gisolfi CV, Duchman SM. Guidelines for optimal replacement beverages for different athletic events. *Med Sci Sports Exerc* 1992;**24**:679–87.
- 12 Almond CSD, Shin AY, Fortescue EB, *et al*. Hyponatremia among runners in the Boston marathon. *N Engl J Med* 2005;**352**:1550–6.
- 13 Convertino VA, Armstrong LE, Coyle EF, *et al*. American College of Sports Medicine Position Stand: exercise and fluid replacement. *Med Sci Sports Exerc* 1996;**28**:i–vii.

The aetiology of exercise-associated hyponatraemia is established and is not “mythical”

Epstein and Cohen-Sivan’s conclusion that the aetiology of exercise-associated hyponatraemia (EAH) is more complex than our “oversimplified” explanation stems from their misunderstanding of how fluid balance is regulated during and after exercise and their ignorance of the conclusions of our most recent publication.¹

In contrast with their assertion that “body fluid homeostasis is a primary variable which is important in the stability of the ‘internal